



Charles J. Aramburo, MD · Kelley S. Dempsey, MD
Buckminster J. Farrow, MD · John D. Fisher, MD
Robert M. Hillery, MD · Thomas Wylie Moore, MD
Guillermo Ponce de Leon, MD · Michael Todd Wood, MD

Post Operative instructions- Gallbladder removal (Cholecystectomy)

Call and schedule your post-op visit to occur approximately 7-10 days after surgery.

Wound care:

- 1) Remove outer dressing (if you have them) at 24-48 hours post-op after surgery.
- 2) Leave the steri-strips in place if you have them. They will fall off on their own.
- 3) You may have skin glue sealing your incision; this resembles super glue and is used to reinforce your dissolvable sutures making your incision waterproof and allowing you to take a shower. Do not soak in a tub or go swimming for 14 days.
- 4) You may use an ice bag to reduce swelling at the surgery site for the first 24 hours and then switch to heat if needed.
- 5) You are able to take a shower 24 hours after surgery. Dry off the surgical incision after showering and cover with dry gauze if needed.

Diet:

You may advance your diet as tolerated after surgery; you should reintroduce fatty and spicy foods slowly, in small portions and monitor how you tolerate them. If you experience nausea or indigestion please resume a bland diet and give your digestive system more time to adjust.

Activity:

For 6 weeks after surgery

- 1) No heavy lifting over 20 lbs.
- 2) No strenuous exercise such as push-ups, sit-ups or abdominal straining.
- 3) Walking is encouraged.
- 4) Normal daily activities are encouraged.
- 5) Running and biking are allowed when you feel up to it.
- 6) No driving until you are pain free and no longer using your pain medication.

Constipation and bloating:

- 1) If you feel bloated, increase your activity.
- 2) If you are constipated, increase your water intake and use an over-the-counter stool softener such as mineral oil or a laxative such as Milk of Magnesia as directed on the bottle.

Call our office if:

- 1) You develop any unusual signs or symptoms such as tenderness, bleeding, redness, or drainage at the surgery site.
- 2) You have a temperature of 100.5 degrees or higher.
- 3) You have yellowing of the skin or eyes.
- 4) You have shortness of breath upon exertion.

Medication refills:

Please call your pharmacy directly and request any needed refills. The pharmacy will contact our office and refills will be expedited.

Memorial Hermann Sugar Land Location:
Methodist Sugar Land Location:
Memorial Hermann Southwest Location:
Memorial Hermann Katy Location:
Central Business Office:

17510 W. Grand Parkway S., Suite 490 · Sugar Land, Texas 77479 · 281.342.8932
16651 Southwest Freeway, Suite 360 · Sugar Land, Texas 77479 · 281.491.6200
7737 Southwest Freeway, Suite 100 · Houston, Texas · 713.772.1200
23920 Katy Freeway, Suite 560A · Katy, Texas · 281.693.3500
10101 Southwest Freeway, Suite 105 · Houston, Texas · 713.255.6300